

The Calculation of Critical Parameters in SU(2) Gauge Theory with Kouvel-Fisher Method

O.A. Mogilevsky

Bogolyubov Institute for Theoretical Physics, 14-B Metrologichna Str., Kiev 03143, Ukraine

Abstract

We calculate the critical coupling $4/g_c^2$ and critical exponent β for the order parameter in SU(2) lattice gauge theory by applying of the finite size scaling technique and the method proposed by Kouvel and Fisher for analysis of experimental data. In contrast to the standard finite size scaling approach, this method allows to determine simultaneously both $g_c^2/4$ and β as two parameters of the linear fit to the Monte-Carlo data.

Lattice calculations by Monte-Carlo (MC) method have so far been the only method for studying the temperature phase transition in gauge theories from the first principles. By their nature, MC calculations are carried out on the finite lattices. This is a most serious drawback of a computer simulations, because no finite system can exhibit a true phase transition. Nevertheless, finite systems remind of phase transitions and systematic studies of these pseudo-transitions as function of system size may reveal information about the phase transition in the thermodynamic limit. One way to do this particularly in order to evaluate the critical exponents at the theory is to use finite size scaling (FSS).

FSS was proposed in statistical physics [1, 2] and applied to high precision MC data of Ising [3, 4] and other models. The validity of this method for gauge theories has been demonstrated through the investigation of critical properties of SU(2) lattice gauge theories at finite temperature [5–9].

In this letter we propose the method of analysis of MC data which is the combination of FSS technique and so-called Kouvel-Fisher method for analysis of experimental data in solid state physics [11]. We consider MC data for SU(2) order parameter on the lattices $8^3 \times 4$, $12^3 \times 4$ and $18^3 \times 4$. In contrast to the standard FSS approach, our method allows to determine the critical coupling $4/g_c^2$ and critical exponent β as two parameters of the linear fit.

The partition function for SU(2) gauge theory on $N_\sigma^3 \times N_\tau$ lattice is

$$Z = \int \prod_{(\mu,\nu)} dU_{\mu,\nu} e^{-S(U)}, \quad (1)$$

where $U_{\mu,\nu}$ are the SU(2) matrices on the link (μ, ν) , S is the standard Wilson action

$$S(U) = \frac{4}{g^2} \sum_p \left(1 - \frac{1}{2} \text{Tr} U_p \right). \quad (2)$$

U_p is a product of link matrices around a plaquette. The temperature T (volume V) is defined by size of the lattice in the time-like (space-like) direction (a is a lattice spacing)

$$T = 1/(N_\tau a), \quad V = (N_\sigma a)^3. \quad (3)$$

The order parameter of the deconfinement phase transition on an infinite volume lattice is the expectation value of lattice average of the Polyakov loop

$$\langle L \rangle = \left\langle \frac{1}{N_\sigma^3} \sum_{\mathbf{n}} L_{\mathbf{n}} \right\rangle, \quad (4)$$

where

$$L_{\mathbf{n}} = \frac{1}{2} \text{Tr} \prod_{\tau=1}^{N_\tau} U_{\tau, \mathbf{n}; 0} \quad (5)$$

and $U_{\tau, \mathbf{n}; 0}$ is the SU(2) link matrix at the point $(\mathbf{n}, \tau)$ in the time-like direction. On the finite lattice one considers as the order parameter $\langle |L| \rangle$, since in this case $\langle L \rangle$ is always zero due to system flips between the two ordered states.

According to FSS theory the scaling function for the order parameter on the lattice $N_\sigma^3 \times N_\tau$, N_τ fix is given by [2]

$$N_\sigma^{\frac{\beta}{\nu}} \langle |L| \rangle = Q_L(x, g_1 N_\sigma^{-y_1}), \quad (6)$$

where β, ν are the critical exponents for order parameter and correlation length correspondingly, g_1 is the irrelevant scaling field, $y_1 \approx 1$. The scaling variable x is defined as

$$x = t \cdot N_\sigma^{\frac{1}{\nu}}, \quad t = \frac{g_c^2/4 - g^2/4}{g_c^2/4}. \quad (7)$$

Our definition of reduce temperature t differs from one used in [7–10]

$$t = \frac{4/g^2 - 4/g_c^2}{4/g_c^2} \quad (8)$$

and seems to be more natural because the coupling constant $g^2/4$ plays just the role of a "temperature" of the effective theory which can be obtained from the partition function (1) by integration on all link variables except those for Polyakov loops. It is clear that this difference disappears in the limit $g^2/4 \rightarrow g_c^2/4$.

At large N_σ the scaling function Q_L must behave as

$$Q_L \sim (x)^\beta. \quad (9)$$

This behaviour may be observed only if x becomes "sufficiently large". Deviations from (9) are in close vicinity of the critical point due to finite size rounding and further away from $x = 0$ due to correction to scaling. Expanding Q_L in (6) around $x = 0$ at large N_σ results in

$$N_\sigma^{\frac{\beta}{\nu}} \langle |L| \rangle = a + \frac{b}{N_\sigma^{y_1}} + \left(c + \frac{d}{N_\sigma^{y_1}} \right) x, \quad (10)$$

where a, b, c, d are unknown constants. Dropping the term $O(N_\sigma^{-y_1})$ one can obtain

$$N_\sigma^{\frac{\beta}{\nu}} \langle |L| \rangle = a + cx, \quad (11)$$

what is exactly the approach of Refs. [7–10]. On the other hand at x "sufficiently large" the x -dependence is drastically changed. In Ref. [6] the scaling function Q_L has been calculated in the form

$$Q_L = Ax^\beta \left[1 + \frac{B}{N_\sigma^{y_1}} x^{y_1 \nu} \right] \quad (12)$$

by the data collapsing method. It has been shown that the correction to scaling term in (12) is negligibly small as compared to the leading term on the interval $0,5 < x < 2$. In both cases the SU(2) Monte-Carlo data for the order parameter have been used which were computed on $N_\sigma^3 \times N_\tau$ lattices with $N_\sigma=8,12,18,26$ and $N_\tau = 4$ and have already reported on in Refs. [5–10]. The numerical value of the critical exponent β has been found which is compatible with the corresponding value $\beta = 0,327$ of the 3- d Ising model [4].

These two procedures have two disadvantages: i) $3-d$ Ising model value of $\nu = 0,631$ has to be used; ii) the very accurate calculation of critical coupling $g_c^2/4$ is necessary. The latter is determining by the fourth cumulant of $\langle|L|\rangle$ as in Refs. [5, 7] or by the χ^2 -method as proposed in Ref. [8]. Even small shift leads to the additional uncertainty in the determination of critical exponent β , because it depends very sensitively on the value inserted for $g_c^2/4$.

We propose below the method for the determination of the critical exponent β which does not require to know of $g_c^2/4$ from the very beginning. No information about the critical exponent ν is also required. Following the Kouvel-Fisher idea which was used for analysis of the experimental data in solid state physics [11] we consider the function

$$K(x) = \frac{Q_L}{\partial Q_L / \partial x}. \quad (13)$$

According to the results of Ref. [6] in the region $0,5 < x < 2$ it is easy to find from (12)

$$K(x) = \frac{x}{\beta} \left[1 + \frac{\text{const}}{N_\sigma^{y_1}} x^{y_1 \nu} \right]. \quad (14)$$

Dropping the correction to scaling term we have in this region with the same accuracy as in (11)

$$K(x) = \frac{x}{\beta}. \quad (15)$$

Taking into account the identity

$$\frac{\partial Q_L}{\partial x} = -\frac{g_c^2}{4} N_\sigma^{-\frac{(1-\beta)}{\nu}} \frac{\partial \langle|L|\rangle}{\partial (g^2/4)} \quad (16)$$

we have immediately

$$K(x) = \frac{x}{\beta} = -\frac{N_\sigma^{\frac{1}{\nu}}}{g_c^2/4} \frac{\langle|L|\rangle}{\partial \langle|L|\rangle / \partial (g^2/4)}. \quad (17)$$

Finally, we obtain the simple but informative expression

$$\frac{\langle|L|\rangle}{\partial \langle|L|\rangle / \partial (g^2/4)} = \frac{1}{\beta} \left(\frac{g^2}{4} - \frac{g_c^2}{4} \right). \quad (18)$$

Eq. (18) means that for the lattices with N_τ fix and various N_σ there are the intervals of $g^2/4$ where MC data for different N_σ must arrange around

the same straight line. The inverse slop of this line represents the critical exponent β and the intersection with $g^2/4$ axis yields the critical coupling.

In order to verify this approach we have used the high precision MC data produced by Bielefeld group on the lattices $8^3 \times 4$, $12^3 \times 4$ and $18^3 \times 4$ [5, 10]. The calculation of derivatives with respect to $g^2/4$ has been made by using the parabolic fit for $\langle |L| \rangle$ (every parabola includes five MC points). This fit does not smooth the statistical fluctuations, but connects the data continuously and with continuous first and second derivatives. The least minimal χ^2 -fit of Eq. (18) ($\chi^2/N = 0,62$) includes 7 MC points in the region $0.396 < g^2/4 < 0.426$ ($2.345 < 4/g^2 < 2.527$) for $N_\sigma = 8$, 24 MC points in the region $0.414 < g^2/4 < 0.430$ ($2.325 < 4/g^2 < 2.415$) for $N_\sigma = 12$ and 16 MC points in the region $0.427 < g^2/4 < 0.433$ ($2.310 < 4/g^2 < 2.340$) for $N_\sigma = 18$. The result is

$$\beta = 0.326 \pm 0.003; \quad 4/g_c^2 = 2.2988 \pm 0.0007. \quad (19)$$

We see that the critical exponent β is in excellent agreement with $3-d$ Ising model value and the inverse critical coupling $4/g_c^2$ coincides with that value which has been calculated in Refs. [7–10] by other method.

In conclusion the following comments are in order. We have shown that the Kouvel-Fisher modification of FSS gives the extremely useful method for the calculation of critical exponent β in SU(2) lattice gauge theory. This method allows to investigate the finite lattice system at small but finite $t = (g^2/4 - g_c^2/4)/(g_c^2/4)$ and large N_σ with infinite volume formulae (18). The critical coupling $g_c^2/4$ and critical exponent β are simultaneously determined as two parameters of the linear fit. It is clear that this approach may be applied to the calculation of all critical exponents in SU(2) gauge theory and to other models which undergo the second-order phase transition.

References

- [1] K. Binder, Z. Phys. **B43** (1981) 119.
- [2] M.N. Barber, R.B. Pearson, D. Toussaint, and J. Richardson, Phys. Rev. **B32** (1985) 1720.
- [3] A.M. Ferrenberg and D.P. Landau, Phys. Rev. **B44** (1991) 5081.

- [4] M. Caselle and M. Hasenbusch, J. Phys. A **30** (1997) 4963; Nucl. Phys. B (Proc. Suppl.) **63** (1998) 613.
- [5] J. Engels, J. Fingberg, and M. Weber, Nucl. Phys. **B332** (1990) 737.
- [6] V.A. Averchenkov and O.A. Mogilevsky, Acta Phys. Slov. **41** (1991) 102.
- [7] J. Engels, J. Fingberg, and D.E. Miller, Nucl. Phys. **B387** (1992) 501.
- [8] J. Engels, J. Fingberg, and V.K. Mitrjushkin, Phys. Lett. **B298** (1993) 154.
- [9] J. Engels, S. Mashkevich, T. Scheideler, and G. Zinovjev, Phys. Lett. **B365** (1996) 219.
- [10] J. Engels and T. Scheideler, Nucl. Phys. **B539** (1999) 557.
- [11] J.S. Kouvel and M.S. Fisher, Phys. Rev. **A136** (1964) 1626.
P. Braun and M. Föhnle, Journ. Stat. Phys. **52** (1988) 775.